

Gradient estimates for a nonlinear parabolic equation under Ricci flow

Shu-Yu Hsu

Department of Mathematics
National Chung Cheng University
168 University Road, Min-Hsiung
Chia-Yi 621, Taiwan, R.O.C.
e-mail:syhsu@math.ccu.edu.tw

June 25, 2008

Abstract

Let $(M, g(t))$, $0 \leq t \leq T$, be a n -dimensional complete noncompact manifold, $n \geq 2$, with bounded curvatures and metric $g(t)$ evolving by the Ricci flow $\frac{\partial g_{ij}}{\partial t} = -2R_{ij}$. We will extend the result of L. Ma and Y. Yang and prove a local gradient estimate for positive solutions of the nonlinear parabolic equation $\frac{\partial u}{\partial t} = \Delta u - au \log u - qu$ where $a \in \mathbb{R}$ is a constant and q is a smooth function on $M \times [0, T]$.

Key words: complete noncompact manifold, Ricci flow, local gradient estimate, nonlinear parabolic equation

Mathematics Subject Classification: Primary 58J05, 58J35

1 Introduction

Recently there is a lot of studies on Ricci flow on manifolds by R. Hamilton and others [6], [5], because Ricci flow is a powerful tool in analyzing the structure of manifolds. Let $(M, g(t))$, $0 \leq t \leq T$, be a n -dimensional complete noncompact manifold with metric $g(t)$ evolving by the Ricci flow

$$\frac{\partial g_{ij}}{\partial t} = -2R_{ij} \quad \text{in } M. \quad (1.1)$$

$(M, g(t))$ is said to be a gradient Ricci soliton ([6],[12]) if there exists a constant $c \in \mathbb{R}$ such that

$$\text{Ric} = cg + D^2f \quad \text{in } M \quad (1.2)$$

where D^2f is the Hessian of f . $(M, g(t))$ is a expanding gradient soliton if $c < 0$ and $(M, g(t))$ is a shrinking gradient soliton if $c > 0$. As observed by L. Ma [10] if we let $f = \log u$, then after some computation using (1.2) we get

$$\Delta u + 2cu \log u = (A_0 - nc)u \quad \text{in } M \quad (1.3)$$

for some constant A_0 . Then the study of the gradient soliton on M is reduced to the study of the properties of the solution u of (1.3). In [10] L. Ma using the technique of S.Y. Cheng, P. Li and S.T. Yau [4], [9], proved the local gradient estimates of positive solutions of the equation

$$\Delta u - au \log u - bu = 0 \quad \text{in } M$$

where $a > 0$ and $b \in \mathbb{R}$ are constants for complete noncompact manifolds with a fixed metric and curvature locally bounded below. In [13] Y. Yang extended L. Ma's result and proved the local gradient estimates of positive solutions of the equation

$$u_t = \Delta u - au \log u - bu \quad \text{in } M \times (0, T]$$

where $a, b \in \mathbb{R}$ are constants for complete noncompact manifolds with a fixed metric and curvature locally bounded below. In [3] A. Chau, L.F. Tam and C. Yu, proved the local gradient estimates of positive solutions of the conjugate heat equation

$$u_t = \Delta^t u - qu \quad \text{in } M \times (0, T]$$

where q is a smooth function on $M \times [0, T]$ and Δ^t is the Laplace operator with respect to the metric $g(t)$ for any complete noncompact manifold $(M, g(t))$, $0 \leq t \leq T$, with bounded curvature and metric g satisfying

$$\frac{\partial g_{ij}}{\partial t} = 2h_{ij} \quad \text{in } M \times (0, T] \quad (1.4)$$

where $h(x, t) = (h_{ij}(x, t))$ is a smooth symmetric tensor on $M \times [0, T]$. In this paper we will extend their results and prove the local gradient estimates of positive solutions of the equation

$$u_t = \Delta^t u - au \log u - qu \quad \text{in } M \times (0, T] \quad (1.5)$$

where $a \in \mathbb{R}$ is a constant and q is a smooth functions on $M \times [0, T]$ for any complete noncompact manifold $(M, g(t))$, $0 \leq t \leq T$, with bounded curvature and metric g satisfying (1.4). In particular our result holds for complete noncompact manifolds with metric evolving by Ricci flow.

We start with some definitions. Let ∇^t and Δ^t be the covariant derivative and Laplacian with respect to the metric $g(t)$. When there is no ambiguity, we will drop

the superscript and write ∇ , Δ , for ∇^t , Δ^t , respectively. For any $r > 0$, $x_0 \in M$, $0 < t_1 \leq t_0 \leq T$, let $B_r(x_0)$ be the geodesic ball with center x_0 and radius r with respect to the metric $g(0)$. For any $x_1, x_2 \in M$, let $r(x_1, x_2)$ be the distance between x_1 and x_2 with respect to the metric $g(0)$.

2 Local gradient estimates

In this section we will prove the local gradient estimates for positive solutions of (1.5). We will let $(M, g(t))$, $0 \leq t \leq T$, be a complete noncompact manifold with metric g satisfying (1.4) and

$$|Rm| \leq k_0 \quad \text{on } M \times [0, T]$$

for some constant $k_0 > 0$ for the rest of the paper. Similar to [3] we will assume that q is a smooth functions on $M \times [0, T]$ and $h = (h_{ij})$ is a smooth symmetric tensor on $M \times [0, T]$. We will also assume the following for the metric g and symmetric tensor $h(x, t) = (h_{ij}(x, t))$:

(A1) $\|h\|, \|\nabla^t h\|$ are uniformly bounded on $[0, T]$ where the norms are taken on M .

(A2) $|q|, |\nabla^t q|, |\Delta q|$ are uniformly bounded on $M \times [0, T]$.

We also let $H = g^{ij}h_{ij}$ be the trace of $h = (h_{ij})$.

Theorem 2.1. *Let u be a positive solution of (1.5). Then for any $\alpha > 1$ and $0 < \delta < 1$ there exists a constant $C_1 > 0$ depending on k_0 , α , δ , $a_+ = \max(a, 0)$ and the space-time uniform bound of $\|h\|, \|\nabla^t h\|, |q|, |\nabla^t q|, |\Delta^t q|$ such that*

$$\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} - \alpha a \log u - \alpha q \leq \frac{n\alpha^2}{2(1-\delta)t}(1 + ta_-) + C_1 \quad \text{in } M \times (0, T] \quad (2.1)$$

where $a_- = -\min(a, 0)$.

Proof: We will use a modification of the technique of [3] and [8] to prove the theorem. Let $f = \log u$. Then by (1.5) f satisfies

$$f_t = \Delta f + |\nabla f|^2 - af - q. \quad (2.2)$$

By [9] and [3] f satisfies

$$\begin{cases} (\Delta f)_t = \Delta f_t - 2h_{ij}f_{ij} - 2h_{ik;i}f_k + \nabla H \cdot \nabla f \\ (|\nabla f|^2)_t = 2\nabla f_t \cdot \nabla f - 2h(\nabla f, \nabla f) \end{cases} \quad (2.3)$$

where $h(\nabla f, \nabla f) = h_{ij}f_i f_j$. Let

$$F(x, t) = t[|\nabla f|^2 - \alpha(f_t + af + q)]. \quad (2.4)$$

By (2.3) and the Weitzenbock-Bochner formula [1],

$$\Delta(|\nabla f|^2) = 2\nabla f \cdot \nabla(\Delta f) + 2|\nabla^2 f|^2 + 2R_{ij}f_i f_j,$$

we have in normal coordinates

$$\begin{aligned} t^{-1}\Delta F &= \Delta(|\nabla f|^2) - \alpha\Delta f_t - \alpha a\Delta f - \alpha\Delta q \\ &= 2|\nabla^2 f|^2 + 2\nabla f \cdot \nabla(\Delta f) + 2R_{ij}f_i f_j - \alpha(\Delta f)_t - 2\alpha h_{ij}f_{ij} - 2\alpha h_{ik;i}f_k \\ &\quad + \alpha\nabla H \cdot \nabla f - \alpha a\Delta f - \alpha\Delta q. \end{aligned} \quad (2.5)$$

Now by (2.2) and (2.4),

$$\Delta f = f_t + af + q - |\nabla f|^2 = \frac{1}{\alpha} \left(|\nabla f|^2 - \frac{F}{t} \right) - |\nabla f|^2 = -\frac{1}{\alpha} \left((\alpha - 1)|\nabla f|^2 + \frac{F}{t} \right). \quad (2.6)$$

Then by (1.4), (2.2), (2.4) and (2.6),

$$\begin{aligned} & -\alpha(\Delta f)_t + 2\nabla f \cdot \nabla(\Delta f) \\ &= \frac{\partial}{\partial t} \left(\frac{F}{t} + (\alpha - 1)|\nabla f|^2 \right) + 2\nabla f \cdot \nabla(f_t + af + q - |\nabla f|^2) \\ &= \frac{F_t}{t} - \frac{F}{t^2} + 2\alpha\nabla f \cdot \nabla f_t + 2\nabla f \cdot \nabla(af + q - |\nabla f|^2) - 2(\alpha - 1)h_{ij}f_i f_j \\ &= \frac{F_t}{t} - \frac{F}{t^2} + 2\alpha\nabla f \cdot \nabla \left(\frac{1}{\alpha}|\nabla f|^2 - \frac{F}{\alpha t} - af - q \right) + 2\nabla f \cdot \nabla(af + q - |\nabla f|^2) \\ &\quad - 2(\alpha - 1)h_{ij}f_i f_j \\ &= \frac{F_t}{t} - \frac{F}{t^2} - \frac{2}{t}\nabla f \cdot \nabla F - 2(\alpha - 1)a|\nabla f|^2 - 2(\alpha - 1)\nabla f \cdot \nabla q - 2(\alpha - 1)h_{ij}f_i f_j. \end{aligned} \quad (2.7)$$

Hence by (2.5), (2.6) and (2.7),

$$\begin{aligned} & t^{-1}(\Delta F - F_t + 2\nabla f \cdot \nabla F) \\ &= 2 \sum_{i,j} (f_{ij}^2 - \alpha h_{ij}f_{ij}) + 2R_{ij}f_i f_j - (\alpha - 1)a|\nabla f|^2 - 2(\alpha - 1)\nabla f \cdot \nabla q \\ &\quad - 2(\alpha - 1)h_{ij}f_i f_j - \frac{F}{t^2} + a\frac{F}{t} - 2\alpha h_{ik;i}f_k + \alpha\nabla H \cdot \nabla f - \alpha\Delta q. \end{aligned} \quad (2.8)$$

By (2.8) and Young's inequality, for any $0 < \delta < 1$ there exist constants $C_1 > 0$, $C_2 > 0$, $C_3 > 0$ and $C_4 > 0$ which depend on a_+ and such that

$$\begin{aligned} & t^{-1}(\Delta F - F_t + 2\nabla f \cdot \nabla F) \\ &\geq 2(1 - \delta) \sum_{i,j} f_{ij}^2 - C_1(|\nabla f| + |\nabla f|^2) - C_2 - \frac{F}{t^2} + a\frac{F}{t} \\ &\geq \frac{2(1 - \delta)}{n}(\Delta f)^2 - C_3|\nabla f|^2 - C_4 - \frac{F}{t^2} + a\frac{F}{t}. \end{aligned} \quad (2.9)$$

By (1.4) and (A1) there exist constants $c_1 > 0$, $c_2 > 0$, such that

$$\begin{cases} c_1 g_{ij}(x, 0) \leq g_{ij}(x, t) \leq c_2 g_{ij}(x, 0) & \forall x \in M, 0 \leq t \leq T \\ c_1 g^{ij}(x, 0) \leq g^{ij}(x, t) \leq c_2 g^{ij}(x, 0) & \forall x \in M, 0 \leq t \leq T. \end{cases} \quad (2.10)$$

Let $x_0 \in M$ and $r(x) = r(x, x_0)$. Then by (2.10) there exists a constant $c_3 > 0$ such that

$$|\nabla^t r(x)|^2 \leq c_3 \quad \text{in } M \quad \forall 0 \leq t \leq T. \quad (2.11)$$

Let $\psi \in C^\infty([0, \infty))$, $0 \leq \psi \leq 1$, be such that $\psi(r) = 1$ for all $0 \leq r \leq 1$, $\psi(r) = 0$ for all $r \geq 2$ and $\psi'' \geq -c_4$, $0 \leq \psi' \leq c_4$ and $\psi'^2/\psi \leq c_4$ on $[0, \infty)$ for some constant $c_4 > 0$. Let $R \geq 1$ and

$$\phi(x) = \psi\left(\frac{r(x)}{R}\right).$$

Then

$$\frac{|\nabla \phi|^2}{\phi} = \frac{|\nabla^t r(x)|^2}{R^2} \cdot \frac{\psi'^2}{\psi} \leq \frac{c_3 c_4}{R^2}. \quad (2.12)$$

By an argument similar to [2] and [9] we may assume without loss of generality that $\phi(x)$ and $r(x)$ are both smooth on M . By (2.10), (2.11), the Laplacian comparison theorem [11] and an argument similar to the proof of Lemma 1.3 of [7] there exists a constant $c_5 > 0$ such that

$$\begin{aligned} \Delta^t r(x) &\leq c_5 \quad \forall x \in M, r(x) \geq R, 0 \leq t \leq T \\ \Rightarrow \quad \Delta^t \phi &= \psi'' \frac{|\nabla^t r(x)|^2}{R^2} + \psi' \frac{\Delta^t r(x)}{R} \geq -\frac{c_3 c_4}{R^2} - \frac{c_4 c_5}{R} \geq -C_6(R^{-1} + R^{-2}) \end{aligned} \quad (2.13)$$

in $M \times [0, T]$ for some constant $C_6 > 0$. Suppose the function ϕF attains its maximum at (x_0, t_0) . If $(\phi F)(x_0, t_0) \leq 0$ for any $R \geq 1$, then (2.1) holds in $M \times [0, T]$ and we are done. Hence we may assume without loss of generality that there exists $R_1 \geq 1$ such that for any $R \geq R_1$, $(\phi F)(x_0, t_0) > 0$. Then at (x_0, t_0) , $F_t \geq 0$,

$$\nabla(\phi F) = 0 \quad \Rightarrow \quad \phi \nabla F = -F \nabla \phi \quad (2.14)$$

and by (2.2), (2.9), (2.12), (2.13) and (2.14),

$$\begin{aligned} 0 &\geq \Delta(\phi F) \\ &\geq \phi \Delta F + 2 \nabla \phi \cdot \nabla F + F \Delta \phi \\ &\geq t_0 \phi \frac{2(1-\delta)}{n} (\Delta f)^2 - C_3 t_0 \phi |\nabla f|^2 - C_4 t_0 \phi - \frac{F}{t_0} \phi + a \phi F + \phi F_t + 2 F \nabla f \cdot \nabla \phi \\ &\quad - 2 F \frac{|\nabla \phi|^2}{\phi} - C_6 F (R^{-1} + R^{-2}) \\ &\geq t_0 \phi \frac{2(1-\delta)}{n} (|\nabla f|^2 - (f_t + a f + q))^2 - C_3 t_0 \phi |\nabla f|^2 - C_4 t_0 \phi - \frac{F}{t_0} \phi - a_- \phi F \\ &\quad - C_5 F |\nabla f| \phi^{\frac{1}{2}} R^{-1} - C_6 F (R^{-1} + R^{-2}) \end{aligned} \quad (2.15)$$

for some constant $C_5 > 0$. Let

$$\sigma = \frac{\alpha^2 - 1}{\alpha^2}, \quad \delta_1 = \frac{2(\alpha - 1)}{\alpha^2} \quad \text{and} \quad \tau = \frac{1}{2}(\sigma - \delta_1).$$

Then by (2.15) and an argument similar to the proof on P.13–14 of [3],

$$\begin{aligned} 0 &\geq -\phi F[C_7 t_0(1 + \delta_1^{-1})(R^{-1} + R^{-2}) + 1 + t_0 a_-] - C_8(1 + \tau^{-1})t_0^2 \\ &\quad + \frac{2(1 - \delta)}{n}(1 - \sigma)(t_0 \phi)^2(|\nabla f|^2 - \alpha(f_t + af + q))^2 \\ &\geq -\phi F[C_7 t_0(1 + \delta_1^{-1})(R^{-1} + R^{-2}) + 1 + t_0 a_-] - C_8(1 + \tau^{-1})t_0^2 + \frac{2(1 - \delta)}{n}\alpha^{-2}(\phi F)^2 \end{aligned} \quad (2.16)$$

for some constant $C_7 > 0$ and $C_8 > 0$. Let $y = \phi F$ and $b = C_7 t_0(1 + \delta_1^{-1})(R^{-1} + R^{-2}) + 1 + t_0 a_-$. Then by (2.16),

$$\begin{aligned} y^2 - \frac{n\alpha^2 b}{2(1 - \delta)}y &\leq \frac{C_8 n}{2(1 - \delta)}\alpha^2(1 + \tau^{-1})t_0^2 \\ \Rightarrow \left(y - \frac{n\alpha^2 b}{4(1 - \delta)}\right)^2 &\leq \left(\frac{n\alpha^2 b}{4(1 - \delta)}\right)^2 + \frac{C_8 n}{2(1 - \delta)}\alpha^2(1 + \tau^{-1})t_0^2 \\ \Rightarrow y &\leq \frac{n\alpha^2}{2(1 - \delta)}[C_7 T(1 + \delta_1^{-1})(R^{-1} + R^{-2}) + 1 + Ta_-] + C_9 T. \end{aligned} \quad (2.17)$$

where $C_9 = \alpha \left(\frac{C_8 n}{2(1 - \delta)}\right)^{\frac{1}{2}} (1 + \tau^{-1})^{\frac{1}{2}}$. Then for any $x \in B_R(x_0)$,

$$F(x, T) \leq \frac{n\alpha^2}{2(1 - \delta)}[C_7 T(1 + \delta_1^{-1})(R^{-1} + R^{-2}) + 1 + Ta_-] + C_9 T$$

Replacing T by t and letting $t \in (0, T]$ we have for any $x \in B_R(x_0)$,

$$\begin{aligned} |\nabla f|^2 - \alpha f_t - \alpha af - \alpha q &\leq \frac{n\alpha^2}{2(1 - \delta)}[C_7(1 + \delta_1^{-1})(R^{-1} + R^{-2}) + \frac{1}{t} + a_-] + C_9 \\ \Rightarrow |\nabla f|^2 - \alpha f_t - \alpha af - \alpha q &\leq \frac{n\alpha^2}{2(1 - \delta)t}(1 + ta_-) + C_9 \quad \text{as } R \rightarrow \infty. \end{aligned}$$

and the theorem follows. $\square$

Corollary 2.2. *Let $(M, g(t))$, $0 \leq t \leq T$, be a complete noncompact manifold with metric g evolving by the Ricci flow (1.1) and*

$$|\nabla^i Rm| \leq k_0 \quad \text{on } M \times [0, T] \quad \forall i = 0, 1$$

for some constant $k_0 > 0$. Let u be a positive solution of (1.5). Then for any $\alpha > 1$ and $0 < \delta < 1$ there exists a constant $C_1 > 0$ depending on k_0, α, δ, a_+ , the space-time uniform bound of $|q|, |\nabla^t q|, |\Delta^t q|$ such that

$$\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} - \alpha a \log u - \alpha q \leq \frac{n\alpha^2}{2(1-\delta)t}(1 + ta_-) + C_1 \quad \text{in } M \times (0, T].$$

Corollary 2.3. Let (M, g) be a complete noncompact manifold with a fixed metric g satisfying

$$|Rm| \leq k_0 \quad \text{in } M$$

for some constant $k_0 > 0$. Let u be a positive solution of

$$\Delta u - au \log u - qu = 0 \quad \text{in } M$$

where q is a smooth function on M with $|q|, |\nabla q|$ and $|\Delta q|$ uniformly bounded on M . Then there exists a constant $C_2 > 0$ such that

$$u \leq C_2 \quad \text{in } M \quad \text{if } a < 0$$

and there exists a constant $C_3 > 0$ such that

$$u \geq C_3 \quad \text{in } M \quad \text{if } a > 0.$$

Proof: By Theorem 2.1 with $h_{ij} = 0, \alpha = 2, \delta = 1/2$, there exists a constant $C_1 > 0$ such that (2.1) holds for all $t > 0$. Letting $t \rightarrow \infty$ in (2.1),

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - 2a \log u - 2q \leq 4na_- + C_1 \\ \Rightarrow & -2a \log u - 2q \leq 4na_- + C_1 \\ \Rightarrow & \begin{cases} u \leq e^{(4na_- + C_1 + 2\|q\|_\infty)/2|a|} & \text{in } M & \text{if } a < 0 \\ u \geq e^{-(C_1 + 2\|q\|_\infty)/2a} & \text{in } M & \text{if } a > 0 \end{cases} \end{aligned}$$

and the corollary follows. $\square$

References

- [1] T. Aubin, *Nonlinear Analysis on manifolds, Monge-Ampère equations*, Springer-Verlag, New York, 1982.
- [2] E. Calabi, *An extension of E. Hopf's maximum principle with application to Riemannian geometry*, Duke J. Math., **25**, (1957), 45–56.
- [3] A. Chau, L.F. Tam and C. Yu, *Pseudolocality for the Ricci flow and applications*, <http://arxiv.org/abs/math/0701153>.

- [4] S.Y. Cheng and S.T. Yau, *Differential equations on Riemannian manifolds and their geometric applications*, Comm. Pure Appl. Math., **28**, (1975), 333–354.
- [5] B. Chow and P. Lu and Lei Ni, *Hamilton's Ricci flow*, Graduate Studies in Mathematics, vol. 77, Amer. Math. Soc., Providence, R.I., U.S.A., 2006.
- [6] R.S. Hamilton, *The formation of singularities in the Ricci flow*, *Surveys in differential geometry*, Vol. II (Cambridge, MA, 1993), 7–136, International Press, Cambridge, MA, 1995.
- [7] S.Y. Hsu, Uniqueness of solutions of Ricci flow on complete noncompact manifolds, <http://arxiv.org/abs/0704.3468>.
- [8] S.Y. Hsu, *Some results for the Perelman LYH-type inequality*, preprint, <http://arxiv.org/abs/math/0801.3506>.
- [9] P. Li and S.T. Yau, *On the parabolic kernel of the Schrödinger operator*, Acta Math., **156**, (1986), 153–201.
- [10] L. Ma, *Gradient estimates for a simple elliptic equation on complete non-compact Riemannian manifolds*, J. Functional Anal., **241**, (2006), 374–382.
- [11] R. Schoen and S.T. Yau, *Lectures on differential geometry*, in 'Conference proceedings and Lecture Notes in Geometry and Topology 1, International Press, 1994.
- [12] P. Topping, *Lectures on the Ricci flow*, London Math. Soc. Lect. Note Ser., 325, Cambridge Univ. Press, Cambridge, 2006.
- [13] Y. Yang, *Gradient estimates for a nonlinear parabolic equation on Riemannian manifold*, Proceeding of A.M.S., electronically published on June 11, 2008.